

**A new species of the genus *Anthrenus* from Pakistan  
(Coleoptera: Dermestidae: Megatominae: Anthrenini)**

Jiří HÁVA<sup>1)</sup> & Zubair AHMED<sup>2)</sup>

<sup>1)</sup>Department of Forest Protection and Entomology,  
Faculty of Forestry and Wood Sciences,  
Czech University of Life Sciences,  
Kamýcká 1176, CZ-165 21, Prague 6 - Suchbát, Czech Republic  
e-mail: jh.dermestidae@volny.cz

<sup>2)</sup>Department of Zoology, Federal Urdu University of Arts,  
Science and Technology, Karachi, Pakistan

**Taxonomy, new species, Coleoptera, Dermestidae, Megatominae, Anthrenini, *Anthrenus*, Pakistan**

**Abstract.** A new species *Anthrenus* (*Nathrenus*) *narani* sp. nov., from Pakistan is described, illustrated and compared with similar species *A. (N.) verbasci* (Linnaeus, 1767) and *A. (N.) pubifer* Reitter, 1899.

## INTRODUCTION

The genus *Anthrenus* Geoffroy, 1762 recently contains 248 species and subspecies in 10 subgenera. Twelve species have been still known from Pakistan (Hamshi & Tashfeen 1992, Háva 2003, 2007, Herrmann & Háva 2009). A new recently collected species is described below.

## MATERIAL AND METHODS

The size of the beetles or of their body parts can be useful in species recognition and thus, the following measurements were made:

total length (TL) - linear distance from anterior pronotal margin to elytral apex.

elytral width (EW) - maximum linear transverse distance.

The following abbreviation are used in the text:

JHAC Private Entomological Laboratory & Collection, Únětice u Prahy, Prague-west, Czech Republic.

Type specimen was labelled with red, printed label bearing the text as follows: "HOLOTYPE *Anthrenus* (*Nathrenus*) *narani* sp. nov. J. Háva & Z. Ahmed det. 2013".

## TAXONOMY

### Subfamily Megatominae Tribe Anthrenini

#### *Anthrenus (Nathrenus) narani* sp. nov.

(Figs. 1-6)

**Type material.** Holotype (♂): N Pakistan, Khyber-Pakhtunhwa Province, Naran env., Malkandi [small town hilly area 52 km of Naran], 18.vi.2010, Z. Ahmed lgt., (JHAC).

**Description.** Holotype. Body measurements (mm): TL 4.3 EW 2.6; body large, oval, elytra broader behind middle (Figs 1-2). Integument of elytra and pronotum castaneous brown, integument of head dark brown. Dorsal surface covered with black and white scales, ventral side mainly with whitish scales, with some intermixed black ones.

Head with black scales with two large spots of white scales (Fig. 3). Labial palpi entirely brown. Antennae with 11 antennomeres; antennomeres dark brown; antennal club oblong oval, consisting of 3 antennomeres (Fig. 4). Eyes large, with brown microsetae; inner margin not emarginate. Median ocellus present on frons.

Pronotum discally with blackish scales, with two spots of white scales, lateral parts with fasciae bearing whitish scales. Antennal fossa broad, closed and conforming to shape of antennal club. Prosternum covered with whitish scales only.

Elytra covered with blackish and whitish scales; the white scales forming fasciae: the first fascia starts before middle and reaches the shoulders and scutellum; the second fascia is located behind middle; the third one is situated near apex and reaches apex at elytral suture. Other parts covered with blackish scales. Individual scales narrow and long. Epipleuron very short, brown, with whitish scales.

Mesosternum and metasternum covered with whitish scales only.



Visible ventrites covered with whitish scales; ventrites I-IV with small spots of blackish scales at lateral margins. The ventrite V without spots; 9<sup>th</sup> ventrite as in Fig. 5. Pygidium brown with brown setation.

Legs entirely brown, with some short, light brown setae, femora also with few whitish scales.

Male genitalia as in Fig. 6.

**Sexual dimorphism.** Female unknown.

**Differential diagnosis.** Each antenna consists of 11 antennomeres, eyes have a complete median margin and thus, the new species belongs to the subgenus *Nathrenus* Casey, 1900. The new species is extremely large compared to other known *Nathrenus* species. Only one species of the subgenus has been known from Pakistan - *A. (N.) verbasci* (Linnaeus, 1767); new species differs from it by its large body, narrow scales and structure of antennae and male genitalia; from other similar species *A. (N.) pubifer* Reitter, 1899 (Turkey, Iran, Iraq, Syria) it differs by its body, dark brown antennae and structure of male genitalia.



Figs 1-6. *Anthrenus (Nathrenus) narani* sp. nov. (holotype): 1- habitus, dorsal aspect; 2- habitus lateral aspect; 3- head; 4- antenna of male; 5- 9<sup>th</sup> ventrite; 6- male genitalia.

**Etymology.** Named according to the type locality - Naran town.

**ACKNOWLEDGEMENTS.** The research was supported by the Internal Grant Agency (IGA n.20124364), Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague.

## REFERENCES

- HASHMI A. A. & TASHFEEN A. 1992: Coleoptera of Pakistan. *Proceedings of Pakistan Congress of Zoology* 12: 133-170.
- HÁVA J. 2003: World Catalogue of the Dermestidae (Coleoptera). *Studie a Zprávy Oblastního Muzea Praha-východ v Brandýse nad Labem a Staré Boleslavi*, Supplementum 1: 1-196.
- HÁVA J. 2007: Dermestidae: 57, 299-320. In: LÖBL I. & SMETANA A. (eds.): *Catalogue of Palaearctic Coleoptera*. Volume 4. *Elateroidea, Derodontoidea, Bostrichoidea, Lymexyloidea, Cleroidea and Cucujoidea*. Stenstrup, Apollo Books: 935 pp.
- HERRMANN A. & HÁVA J. 2009b: *Anthrenus (Anthrenops) kadeji* sp. n. (Coleoptera: Dermestidae: Anthrenini) from Pakistan. *Studies and Reports of District Museum Prague-East Taxonomical Series* 5: 61-64.

*Received: 11.9.2013*  
*Accepted: 31.10.2013*